POSTER PRESENTATION





Treatment for femoroacetabular impingement: a qualitative method for exploring equipoise amongst hip arthroscopy surgeons

Peter Wall, Alba Realpe^{*}, Damian Griffin, Rachel Hobson, Ann Adams

From 2nd Clinical Trials Methodology Conference: Methodology Matters Edinburgh, UK. 18-19 November 2013

The published literature suggests uncertainty about whether operative or nonoperative treatments are best for femoroacetabular impingement (FAI). Without the same level of uncertainty (equipoise) amongst surgeons, a RCT will be challenging. A qualitative study was conducted to explore the level of equipoise amongst arthroscopic FAI surgeons. In phase 1, 14 hip arthroscopy surgeons were interviewed and asked to make treatment decisions based on real life cases that included actively recruiting patients to a theoretical RCT. In phase 2, 9 hip arthroscopy hip surgeons participating in a pilot RCT were interviewed about their experiences so far of taking part in a pilot RCT. Five surgeons took part in both phase 1 and 2. Sixteen (89%) surgeons believed that they were in equipoise and that a RCT was required to generate superior scientific evidence and guidelines for the care. Despite this 5 (36%) surgeons showed a lack of active clinical equipoise when faced with real life case scenarios or discussing involvement with a pilot RCT. Some of the reasons behind surgeons' lack of equipoise, ranged from lack of belief in the FAI pathology, to personal enthusiasm and gut instinct about the efficacy of surgery on one hand; but conservatism on the other. Although many would like a RCT to guide care, there may be particular challenges amongst this same population when actively recruiting patients to a RCT. Qualitative methodology can be used to help design surgical RCTs and address any subsequent difficulties with recruitment.

Warwick Medical School, University of Warwick, Coventry, UK

Published: 29 November 2013

doi:10.1186/1745-6215-14-S1-P103

Cite this article as: Wall *et al.*: Treatment for femoroacetabular impingement: a qualitative method for exploring equipoise amongst hip arthroscopy surgeons. *Trials* 2013 14(Suppl 1):P103.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

) Bio Med Central

Submit your manuscript at www.biomedcentral.com/submit



© 2013 Wall et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.